

Stool-derived eukaryotic RNA assay approximates disease activity index and predict therapeutic response in patients with Crohn's disease on advanced therapies

Erica K. Barnell^{1,2}, Ryan Ghannam², Ali Osman^{3,4}, Richard Roberts², Patrick Donohough¹, Spencer King², Jack Land², Clayton Grass², Matthew A. Ciorba³, Parakkal Deepak³



¹Department of Medicine, Washington University School of Medicine, St. Louis, MO, USA

²Geneoscopy, Saint Louis MO, USA

³Inflammatory Bowel Diseases Center, Division of Gastroenterology, Washington University School of Medicine, St. Louis, MO, USA;

⁴Division of Hospital Medicine, Washington University School of Medicine, St. Louis, MO, USA


RNA calprotectin levels were more specific than protein calprotectin levels when correlating with disease and disease severity.



Classifiers were 81% accurate in parsing subjects with active disease from those in remission and 92% accurate in parsing subjects with moderate disease from those with mild disease.

IBD OUTPUT REPORT

MEASURING RNA CALPROTECTIN



ASSESSING DISEASE ACTIVITY



IDENTIFYING DYSPLASIA



PREDICTING THERAPEUTIC RESPONSE

	ENTYVIO (vedolizumab)	ITGA4 / ITGB7	SENSITIVE
	STELARA (ustekinumab)	IL12 / IL23	RESISTANT
	REMICADE (infliximab)	TNF / TNFR	RESISTANT

Original Investigation

October 23, 2023

Multitarget Stool RNA Test for Colorectal Cancer Screening


Erica K. Barnell, MD, PhD^{1,2,3}, Elizabeth M. Wurster, PhD², Julie La Rocca, MS³, et al.

Author Affiliations | Article Information


JAMA. 2023;330(16):1760-1768. doi:10.1001/jama.2023.22321

JAMA
The Journal of the American Medical Association

RNA biomarkers showed 94% sensitivity for colorectal cancer and 46% sensitivity for advanced adenomas.



Relative cell abundance and expression of transcripts associated with the therapeutic target permit enrichment of responders.



SAMPLE COLLECTION LOGISTICS

*Geneoscopy can also assess banked frozen stool samples as part of pilot studies and clinical trials

